

# LOAN DOCUMENT TRANSMITTAL

REPORT NUMBER FTD-SD(RS)7-03/3-96

AD NUMBER PD-A067002

THE ATTACHED DOCUMENT HAS BEEN LOANED TO DDC FOR PROCESSING.

THIS COPY IS NOT TO BE MARKED OR MUTI 
LATED. REQUEST THAT SPECIAL HANDLING,

BE PROVIDED IN ORDER THAT THE COPY MAY

BE PROMPTLY RETURNED TO THE LENDER.

THE REPORT SHOULD BE RETURNED TO

## DDC DOCUMENT PROCESSING/DIA-2DIVISION P. Coopa-

ACCESSION FOR		
NTIS GRA		A
DDC TAB		
JUSTIFICATION		
BY	ION/AVAILABILITY CODES	_
A DIST.	AVAIL. AND/OR SPECIAL	

# DO NOT PHOTOGRAPH THIS FORM LOAN DOCUMENT

DDC FORM 70

PREVIOUS EDITIONS ARE OBSOLETE


### FOREIGN TECHNOLOGY DIVISION





CHEMISTRY AND BIOCHEMISTRY OF HYDORCARBONS (SELECTED PORTIONS)





Approved for public release; distribution unlimited.

78 12 2

### **EDITED TRANSLATION**

FTD-ID(RS)T-0313-78

12 April 1978

MICROFICHE NR: 24D - 78-C-000 77/

CHEMISTRY AND BIOCHEMISTRY OF HYDROCARBONS (SELECTED PORTIONS)

English pages: 23

Source: Khimiya i Biokhimiya Uglevodov,

Izd-vo "Nauka", Moscow, 1969, pp. 5-10,

229-232.

Country of origin: USSR

Translated by: Marilyn Olaechea

Requester: FTD/TQIS

Approved for public release; distribution

unlimited.

STR SDR	With Section
MARHORI JUSTIFICA	ICEO CI
87	TION/AVARABILITY 5003
Piernin	
Bist.	AVAIL one/er SPESIAL

THIS TRANSLATION IS A RENDITION OF THE ORIGI-NAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DI-VISION.

PREPARED BY:

TRANSLATION DIVISION FOREIGN TECHNOLOGY DIVISION WP-AFB, OHIO.

#### Table of Contents

U. S. Board on Geographic Names Transliteration System	ii
Table of Contents	1
Intensity of Synthesizing Glycogen From Various Glycogen- Forming Agents with Functional Shifts in Animal Organism,	3.7
by, Z. N. Tupikova, M. I. Prokhorova, et al	17

U. S. BOARD ON GEOGRAPHIC NAMES TRANSLITERATION SYSTEM

Block	Italic	Transliteration	Block	Italic	Transliteration
A a	A a	A, a	Рр	Pp	R, r
Б б	5 6	B, b	Сс	Cc	S, s
Вв	B .	V, v	Тт	T m	T, t.
Гг	Γ :	G, g	Уу	Уу	U, u
Дд	Д д	D, d	ФФ	Φ φ	F, f
Еe	E .	Ye, ye; E, e∗	X ×	XX	Kh, kh
ж ж	ж ж	Zh, zh	Цц	4 4	Ts, ts
3 з	3 ,	Z, z	4 4	4 4	Ch, ch
Ии	и и	I, i	Шш	Шш	Sh, sh
Йй	A a	Y, y	Щщ	Щщ	Shch, shch
Н н	KK	K, k	Ъъ	ъ ъ	n
ת וג	ЛА	L, 1	Ы ы	M w	Y, у
in in	M M	M, m	Ьь	b .	1
Нн.	Н н	N, n	Ээ	9 ,	Е, е
0 0	0 0	0, 0	Юю	10 no	Yu, yu
Пп	Пп	P, p	Яя	ЯЯ	Ya, ya

<sup>\*</sup>ye initially, after vowels, and after ь, ь; e elsewhere. When written as  $\ddot{e}$  in Russian, transliterate as  $y\ddot{e}$  or  $\ddot{e}$ .

#### RUSSIAN AND ENGLISH TRIGONOMETRIC FUNCTIONS

Russian	English	Russian	English	Russian	English
sin	sin	sh	sinh	arc sh	$sinh_{-1}^{-1}$
cos	cos	ch	cosh	arc ch	cosh
tg	tan	th	tanh	arc th	tanh_1
ctg	cot	cth	coth	arc cth	coth_1
sec	sec	sch	sech	arc sch	sech_1
cosec	csc	csch	csch	arc csch	csch <sup>-1</sup>

Russian	English
rot	curl
lg	log

CHEMISTRY AND BIOCHEMISTRY OF HYDROCARBONS

MATERIALS OF THE IV ALL-UNION CONFERENCE ON CHEMISTRY AND BIOCHEMISTRY OF HYDROCARBONS OF 25-31 MAY

TABLE OF CONTENTS

11 Preface

#### MONOSACCHARIDES

13 New group of monosacharides containing sulfur or nitrogen atoms in ring (survey). B.W. Stepanenko.

- Quantum chemistry of hydrocarbons. Yu.A. Zhdanov, V.I.
  Minkin, G.N. Dorofeyenko, Yu.A. Sotroumov, Ye.H. Malysheva.
- 27 Investigation of stereospecific multirotation of monosaccharides. V.A. Pavlov, Ye.I. Klabunovskiy, A.A. Balandin.
- 31 Kinetics and mechanism of acid-base catalysis of certain reatctions of mono- and disaccharides. V.A. Afanas'yev, V.I. Gorykova, V.A. Khaarmats, R.I. Sarybayeva, O.V. Dronov, T.K. Yunusaliyeva, N.I. Trushkina.
- 36 Synthesis and study of structure and properties of N-alkylglycosylamines. B. N. Stepanenko, R.D. Greshnykh.
- Stythesis and study of structure and properties of certain aryl-N-glycosylamines.
- 46 Qualitative and quantitative thin-layer chromatography of N-phenylglycosylamines.
- 48 Radiolysis of aqueous solutions of glucosamine and some of its derivatives.
- 50 Radiolysis of aqueous solutions of glycosides.

- 53 Mass spectrometry of alkylidene derivative polyols. L.S. Golovkina, O.S. Chizhov, N.S. Vul'fson.
- New approaches to synthesizing aminosaccharides. B.a. Dmitriyev, N.E. Bayramova.
- 62 Synthesis of glycosides, oligosaccharides, and regular polysaccharides by the ortho ether method.
- Some reactions of deoxysaccharide halides. K.S. Abanyants.
- 68 Monosaccharide fragments of olivonycins. Yu.A. Berlin.
- 71 Structure of hydrocarbon part of olivomycin antibiotics.
  S. Ye. Yesipov.
- Synthesis of uridine diphospahte-α-D-6-deoxyglucose and its reaction with uridine disphosphate galactosε [UDFG]. N.K. Kochetkov, B.I. Budovskiy, V.N. Shibayev, Yu.Yu. Kusov.
- 78 Investigation of conformation of 2,3-4,5-di-O-isopropylene-β-D- fructose by method of nuclear magnetic

resonance of high resolution. B. I. Glazkov, Ye.M. Malakhayev.

- 81 Effect of structure of pyran cycle on carbocyclization of 1.6-anhydroaldoherose under influence of sodium in liquid ammonia.

  G. V. Davydova, A. V. Lozanova, N. N. Shorygina.
- 84 Chemistry of 1-glycosylindols. M.N. Preobrazhenskaya, M.M. Vigdorchik, L.A. Savel'yeva, N.N. Suvorov.
- 87 On the synthesis of pangamic acid. A.H. Yurdevich, S.G. Verenikina, N.A. Preobrazhenskiy.
- 91 condensation of triphenylmethylene phosphoranes with alforms of monosaccharides. V.G. Alekseyeva, Yu.Ye. Alekseyev.
- 94 Interaction of free aldoses with n-methoxybenzolmethylene phosphorane. B.A. Polenov, Yu.A. Zhdanov.
- 95 a-ketocarbenes in series of hydrocarbons. V.I. Kornilov.

PLANT GLYCOSIDES. CHEMISTRY OF CELLULOSE.

98 New information on the structure of triterpene saponins.

V. Ya. Chirva, V.P. Konyukhov, P.L. Cheban, G.V. Lazur'yevskiy.

100 Securide abiosis - a new biosis from cardiac glycoside - securidaside. V. V. Atula, D.G. Kolesnikov.

103 Enzymatic splitting of acetyl groups in sugar (saccharine)
part of cardiac glycosides. P.I. Gvosdyak.

106 Structure of leontosides A, B, C, D, and E - triterpene glycosides Leontice eversmanii. L.G. Mzhel'skaya, N.K. Abubakirov.

109 On reactivity of hydroxyl groups of cullulose when it interacts with nitrogen oxides. Z.I. Kuznetsova, V.S. Ivanova.

113 Synthesis of 2,3 and 3,6-anhydro derivative celluloses.

L.S. Gal'braykh, Z.A. Rogovin.

116 Problem of interaction of active dyes and cellulose. Ye. V. Rudakova, M.A. Chekalin.

118 Synthesis of C-derivation celluloses with selective replacement of third hydrocarbon atom of glucopyranose chain. R.G. Drylova, S.N. Ryadovskaya, O.P. Golova.

#### CHEMISTRY OF POLYSACCHARIDES (CELLULOSE EXCLUDED).

- 121 General characteristics of polysaccharides of Phaeophyceae.

  V.A. Khomenko.
- 125 Polysaccharides of certain red water plants. A.I. Usov, L.I. Miroshnikova, M.A. Rekhter.
- 127 Polysaccharides of the ginseng family. T.F. Solov'yeva.
- 129 Analysis of monosaccharides of gas-liquid chromatographs.
  Ye.v. Yevtushenko.
- 132 Interation of aqueous solution of Caustic soda and xylan.

  I.S. Kazanskaya, M.S. Dudkin.
- 135 Effect of ammonia on hydrocarbons. N.. Shkant, N.S. Skornyakova, N.A. Lele, A.F. Yatsuk, E.M. Tkhorkhzhevsaya, V.V. Ruzer.
- 138 Comparative study of polysaccharides of leguminous cultures.

  T.Ye. Panasyuk.

139 Polysaccharides of seaweeds (family Zosteraceae). R.G. Ovodova.

CHEMISTRY AND BIOCHEMISTRY OF BIOLOGICALLY ACTIVE POLYSACCHARIDES OF AND ASSORTED BIOPOLYMERS.

141 polysaccharide of cell Walls of diphteritic bacteria. O.K. Orlova.

143 Structure and biological activity of sodium dextrans.

M.Ye. Preobrazhenskaya, V.M. Kuznetsova.

147 Chemical composition and structure of polysaccharides of certain dyed yeast organisms. G. A. Vitovskaya, N.P. Yelinov.

150 Composition, properties and biological activity of prodigiosine. G. Ye. Vaysberg, T.I. Afanas'yeva, L.A. Taranenko, T.N. Likina.

Glycolipids of marine invertebrates. I.G. Zhukova, G.P. Sirnova, V.Ye. Vas'kovskiy, E.Ya. Kostetskiy, T.A. Bogdanovskaya.

- On structure of ribitteykhoic \* acid from actinomycetes producer of the antibiotic auratine. I.B. Naumova. [Translator
  note: "Ribitteykhoic" represents transliteration from Russian. Term
  not positively identified. end note]
- Glycerine teykhoic acid ["teychoic" is transliteration from Russian] of cell wall of Actinomyces antibioticus st. 39. M.Sh. Zaretskaya, I.B. Naumova, A.Z. Shabarova.
- Some data on the composition, properties, and biological activity of lipopolysaccharides of actinomycetes and bacillus coli.

  N.S. Gryaznova.
- 161 Biologically active polysaccharide complex candida tropicalis. Ye.K. Alimova, A.D. Mar'yasina.
- 163 Concentration of fucose in the mucus membrane of the stomach and in blood sermum in presence of ulcer.
- 164 Chemistry of glycoproteins (survey). V.A. Derevitskaya.
- 171 Synthesis and properties of O-glykosides of β-oxy-α-amino

acids. M. G. Bafina, Ye.M. Klyaimov, T.G. Aliyeva.

175 α-L-fucosidase of animal tissues. G.Ya. Vidershayn.

On neuraminic acid and neuraminidase of Myxoviruses during their interaction with sensitive and insensitive cells.

On neuraminidase of diphteritic microbe. Ye.K. Alimova, L.P. Sokolova.

183 Glycolproteid Tamm-Horsfoll under normal conditions and during certain illnesses. N.a. Yusipova, Ye.L. Rozenfel'd.

On action mechanism of lysozyme on chitinoic oligosaccharides. N.A. Kravchenko, Yu.D. Kuznetsov.

191 Carbohydrases of marine invertebrates. Extraction of cellulase Littorina manschurica by gel filtration on biogels in presence of substrate. L.A. Yelyakova, B.V. Sova, E.P. Postinkova, V.Ye. Vas'kovskiy, M.D. martynova, O.S.Chizhov.

GLYCOGEN EXCHANGE

- 195 Mechanism which regulates action of enzymes which participate in glycogen exchange (survey). Ye.L. Rozenfel'd.
- 205 Ribonucleic acid and formation of branching bonds in polysaccharides. A.N. Petrova, R.D. Filippova.
- 208 On certain characteristic features of acid and neutral gamma-amilases of rat liver. I.A. Popova.
- 211 Acid and neutral alpha-oligoglucosidases of animal and human tissues. I.S. Lykomskaya, N.A. Ushakova.
- 215 Extracting and purifying alpha-glucosidase of muscles of while rabbit. N.F. Pisarenko.
- 218 Study of the formation of tetramer, activation and inhibition of phosphorylase b. G.V. Silonova, N.P. Lisovskaya.
- 222 Study of structural and enzymatic properties of phosphorylase b. L.K. Skolysheva, P.L. Bul'fson.
- Regulation of activity and synthesis of glucokinease and glycogensynthesase of the liver. K.I. Shanygina, M.P. Fomina, B.P. Golovin.

- Intensity of synthesizing glycogen from different glycogen-forming agents during functional shifts in living organism.

  Z.N. Tupikova, N.I. Prokhorova, V.A. Vilkova, N.D. Yeshchenko.
- 233 Mechanism by which sugars enter muscle fiber. S.I.

  Vasyanin, N.A. Vinogradova, N.V. Doroshenko, N.N. Nikol'skiy, L.N.

  Pisareva, A.S. Troshin.
- On problem of active center and prosthetic group of PPA of muscles of rabbit. N.D. Dvornikova, N.P. Gulyy, T.T. Volodina, T.N. Pech'nova, N.I. Polikarpova.
- 240 Activity of hexogenase of mormal and denerved skeletal muscles of adult animal and embryo. V.S. Il'yin, V.M. Pleskov, N.I. Razumovskaya.
- Neurohormonal regulatin of activity of glucokinease and utililization of glucose of fat tissue of epoophoron of rats. K.G. Gromova, I.B. Soliternova.
- 248 Change in primary structure and properties of aldonase of skeletal muscles during change in physicological state of organism.

- M.F. Gulyy, L.T. Litvinenko, N.I. Polydarpova, P.P. Martynenko, N.I. Zhuravskiy, V.A. Kokunin.
- 251 Mechanisms which regulate glycolase in early embryogenesis.

  L.S. Mil'man, Yu.G. Yurovitskiy.
- 253 Role of free radical states in conservation of energy in aerobic oxydation of hydrocarbons. L.M. Raykhman.
- 257 Study of adenosine glucose diphosphate and its analogs in various ezymatic reactions. T.N. Druzhinina, N.D. Gabrielyan, A.V. Venkina, M.A. Novikova.
- 260 Effect of nicotine amide on concentration of energy enriched phosphorous compounds in the muscle of the heart under under normal conditions and in the presence of myocarditis. L.A. Tseymlin.
- comparative characteristics of certain properties of dehydrogenase of 3-phosphoroglycerine aldehyde, derived from various sources. S. Ye. Severin, N.K. Nagradova.
- 269 Effect of insulin on activity of dehydrogenase of gucose-6-phosphate and 6-phosphorogluconate of the fatty tissue of epoophoron of normal and fasting rats and rats with alloxan diabetes.

L.S. Uuskyula.

- 272 Some aspects of regulating formation of sedoheptulose in animal tissue. I.D. Golovatskiy.
- Photoinactivation of transketolase. G.A. Kochetov, K.P. Kobylyanskaya.
- 277 On problem of interaction of individual stages of glycolase with pentosophosphate cycle in mucus membrane. I.G. Pupin.

#### HYDROCARBON EXCHANGE IN PATHOLOGICAL STATES

- problem on intercellular regulation of glycolase in skeletal muscle with experimental muscular dystrophy. V.A. Grigor yeva, D.L. Ferdman, L.V. Shchukina.
- On heterogeneity of dehydrogenages of glucoso-6-phosphogluconic acid in heard muscle under normal conditions and in the presence of hypoxia. A.I. Kolotilova, V.K. Vagner, ye.P. Glushankov, V.N. Tkachenko.

286 Effect of glucose and ATF on respiration of ascitic cancer cells

in presence of fluoridase and Ouabian. M.I. Ilinich, N.v. Sarycheva, B.a. Sobchuk, A.V. Genik-Berezovskiy.

288 Glucogenolysis in presence of hyperthermia. I.V.
Vysochina, V.f. Kaverina, I.N. Konstantinova.

290 Concentration of glycogen in brain of rabbits under incluence of psychotropic substances.

Introduction of sulfates of mucopolysaccharides in tissue of cartilage and skin of normal and A-avitaminotic animals. N.N.
Velikiy, R.V. Chagovets.

Nucopolysaccharides of various types of connective tissue in presence of certain pathological states. B.S. Ksavina, N.I, Vinetkaya, T. A. Kol'chinskaya.

Exchange characteristics of gangliosides and cerebrosides of brain in different functional states. S.Yu. Tumanova, M.I. Prokhorova, L.S. Romanova.

HYDROCARBONS AND HYDROCARBON EXCHANGE IN PLANTS

- 301 Hexose phosphates in transport and sugar accumulation in sugar beets. O.A. Pavlinova.
- 305 On problem of mechanisms which regulate activity of invertase in conductive tissues of plants. MV. Turkina, S.V. Sokolova.
- 307 Nucleotide-containing coenzymes of hydrocarbon exchange in green water plants. N.V. Pakhomova, G.N. Zaytseva.
- 309 Simple chromatographic method of fractionating polysaccharides of starch. N.A. Sykova, B.N. Stepanenko.
- 311 Minor component of preparations of plant polysaccharides of Malvaceae family.
- Polysaccharides from stems of hollyhocks and their localization in stem tissues. I.S. Kozhina, N.A. Trukhaleva, N.Ye. Pokina.
- 317 On celluloses of certain types of mould fungi. N.A. Rodionova, N.A. Tiunova, L.I. Martinovich.
- 321 Resolution of the IV Conference on problem of "Chemistry and

Biochemistry of Hydrocarbons."

eo

3

9

1

INTENSITY OF SYNTHESIZING GLYCOGEN FROM VARIOUS GLYCOGEN-FORMING AGENTS WITH FUNCTIONAL SHIFTS IN ANIMAL ORGANISM

Z. N. Tupikova, M. I. Prokhorova, V. A. Vilkova, N. D. Yeshenko

(Physiological Institute im. A. A. Utomskiy of Leningrad University, Laboratory of Biochemistry of Nervous System and Metabolism)

Presented in the present report are materials on a comparative study of the intensity of synthesizing glycogen in the large hemispheres of the brain for various glycogen-forming agents with functional shifts in the organism of the animal.

The study was performed on white rats weighing 100-200 g.

Serving in the capacity of the glycogen predecessor were glucose-1: -C1\* lactate - 1-3-C1\*, and acetate - 1-C1\*, which were introduced subcutaneously at a calculated 10-20 µc per 100 g weight. The length of radioactive exposure was from 10 minutes to 2 hours. Caffine and chloralhydrate were administered subcutaneously at 50 mg per 100 g weight, aminazine - 3-4 mg, insulin - 2 units, adrenalin - 1 mg per 100 g weight of the animal.

Glycogen was isolated from the large hemispheres of the brain by the method of Kerr in the Le Baron modification [6]. The concentration of glycogen was determined by the amount of glucose formed after acid hydrolysis.

The radioactivity of the glycogen was determined by calculating the number of pulses in the glycogen residue; the concentration of lactic acid - by the method of Barker and Summerson [5], its radioactivity - in the residue of the zinc lactate. The radioactivity of the glucose was determined in the glucosazone residue. The number of pulses were calculated from the "Volna" device using end counter TBFL-25.

The obtained results were processed statistically. The values of the mean arithmetic (M) with error (±m) are shown in the tables.

It was established that every 5 min after introduction of the radioactive acetate - 1-C1\* - the specific activity (SA) of the glycogen in the brain constituted a significant value - 106 imp/min/mg. The maximal values of the SA of brain glycogen were found in experiments with 30-60 minutes of exposure. This index decreased somewhat thereafter. It should be mentioned that the curve representing the change in radioactivity of the homogenate in these experiments is more sloping [4]. Comparison of this curve with the curve representing the change in the SA of brain glycogen in time indicates the active entry of the acetate into the glycogen.

The use of marked lactate with a single-hydrocarbon tag (lactate - 1-C1\*) did not make it possible to show a significant amount of it in the brain glycogen [2-4]. Therefore, in the course of further study on the intensity of synthesizing glycogen, we used lactic acid with a solid hydrocarbon tag - lactate - 1-3-C1\*. The studies which were conducted revealed that the maximal SA of brain glycogen in these experiments was found within 60 minutes after the isotope was introduced, when it constituted 121 imp/min/µmole of lactate on the average.

As we see from the results presented in Table 1, glucose is most actively used in glycogen-formation processes in the brain. Within one hour after introduction of the radioactive compound, the SA of

the brain glycogen was more than 600/o of the SA of glucose.

According to our data acetic and lactic acids also participate rather intensively in the synthesis of glycogen in the large hemispheres of the brain, although the use of these compounds in glycogen formation is approximately three times lower than the use of glucose. Based on published data [1, 7, 8] we assume that the synthesis of glycogen in the brain from lactate and, probably, from acetate as well, occurs not as the result of the reversibility of all glycolysis reactions, rather as a result of the routes taken, which guarantee that energy barriers will be overcome-

The administration of chloralhydrate (Table 2) to animals led to a noticeable drop in the SA of brain glycogen, although under these conditions the SA of the predecessor - radioactive lactate - dropped to the same degree. As a result of this the relative specific activity (RSA) of the glycogen remained virtually unchanged. During excitation, which was caused by an injection of caffine, the SA of the lactate in the brain decreased 5.5 times, glycogen - 2.7 times. Thus, the RSA of the glycogen rose to 43.90/o.

The data shown in Table 3 indicate that the SA of the glycogen dropped noticeably as compared to the norm under all effects, while the total activity of the homogenate of the tissue remained almost unchanged (with the exception of experiments where caffine was

introduced, in which case it decreased).

In experiments using glucose-1-6-C14 (see the figure) the intensity of glycogen synthesis in the brain in all effects studied by us was lower than for the norm. The most drastic slowing of glycogen-formation was observed in experiments with adrenalin and insulin, which, as we know, greatly disturb hydrocarbon exchange.

#### BIBLIOGRAPHY

<sup>1.</sup> В. С. Ильин. Механизмы центрального регулирования обмена, их формирование в В. С. Ильин. Механизмы центрального регулирования обмена, их формирование в онтогенезе и нарушения при паталогических состояниях организма. Актовая речь. М., «Наука», 1966.
 М. И. Прохорова, З. Н. Тупикова. Большой практикум по углеводному обмену. Изд. ЛГУ, 1965.
 З. А. Н. Тупикова. В сб. «Нервная система», т. 6, 1965, стр. 11.
 З. Н. Тупикова, В. А. Вилкова, А. М. Корвацкая. Материалы IV Всес. конференции по биохимии нервной системы. Тарту, 1966.
 S. В. Вагкег, W. N. Summerson. J. Biol. Chem., 138, 535, 1941.
 F. N. Le Baron. Biochem. J., 61, 1, 80.
 M. F. Utter. Annual Rev. N. Y. Acad. Sci., 72, 451, 1959.
 M. F. Utter, D. B. Keech, M. C. Scrutton. Advances in Enzyme Regulation, 2, 49, 1964.

	60	р) Удельно ность, тапи	BNAMULE-	
	Радионктивное вещество	(3) saurosen	(4) предше- стеенник	C <sup>14</sup> (OY A).
6770	Глюкоза — 1-6-С14	673±51	1017±46	66,1
??	Ацетат — 1-С14	168±17	720±22°	23,3
v	Лактат — 1-3-С14	658+45	3060+98	21.5

Table 1. Intensity of synthesizing glycogen in brain from various radioactive predecessors. KEY: (1) Radioactive substance, (2) Specific activity, imp/min/g of tissue, (3) Glycogen, (4) Predecessor, (5) Introduction of C14 (relative specific activity), 0/0, (6) Glucose-1-6-C14, (7) Acetate-1-C14, (8) Lactate-1-3-C14, (9) \*In experiments with acetate-C14 the radioactivity of the homogenate of the tissue was used conditionally for the specific activity.

	(1)	Удельная нооть, ил ткани (		(5) УА вликовена	
	Ycaomia onuma	(3) SAUROSEH	Aanmam	VA sakmama ×100	
100000000000000000000000000000000000000	Норма	658±45	3060±98	21,5	
(1)	Хлоралгидрат Кофеин	369±31 243±36	1990±75 553±44	18,5 43,9	

Table 2. Use of radioactive lactate-1-3-C14 for synthesizing glycogen of brain for various functional shifts in animal organisms. KEY: (1) Conditions of experiment, (2) Specific activity, imp/min/g of tissue, (3) Glycogen, (4) Lactate, (5) Specific activity of glycogen/specific activity of lactate, (6) Norm, (7) Chloralhydrate, (8) Caffine.




Fig. 1. Use of glucose-1-6-C1\* for synthesizing glycogen in brain during various functional shifts in animal organisms: 1 - norm, 2 - chloralhydrate, 3 - caffine, 4 - aminazine, 5 - adrenalin, 6 - insulin. KEY: (1) Specific activity of glycogen, % to the norm.

	(n)	Удельная ность, ил тисни (		УА вликовена	
	Yeaoeus onema	(3)	(4)	VA auemama	× 100
(6)	Норма	168±17	720±22	23,3	
(3)	Хлоралгидрат Кофеии	62±12 50±14	624±41 432±35	9,8	
(9)	Инсулик	26±9	708±47	3,7	

Table 3. Use of radioactive acetate-1-C14 for synthesizing brain glycogen during various function shifts in animal organism. KEY: (1) Conditions of experiment, (2) Specific activity, imp/min/g of tissue, (3) Glycogen, (4) Acetate, (5) Specific activity of glycogen/specific activity of acetate, (6) Norm, (7) Chloralhydrate, (8) Caffine, (9) Insulin.

#### DISTRIBUTION LIST

#### DISTRIBUTION DIRECT TO RECIPIENT

ORGAN	IZATION	MICROFICHE	ORGAN	IZATION	MICROFICHE
C509	DMATC DMAAC DIA/RDS-3C USAMIIA BALLISTIC RES LABS AIR MOBILITY R&D	1 2 8 1	E053 E017 E404 E408 E410	AF/INAKA AF/RDXTR-W AEDC AFWL ADTC	1 1 1 1
	LAB/FIO PICATINNY ARSENAL AVIATION SYS COMD	1 1	E413	ESD FTD CCN ASD/FTD/NICD	2 1 3
D008 H300 P005 P055	FSTC MIA REDSTONE NISC USAICE (USAREUR) ERDA CIA/CRS/ADD/SD	5 1 1 1 1		NIA/PHS NICD	1 2
NASA/H AFIT/I		1			

